

Children's Robitussin Elderberry Cough + Chest Congestion DM

Dextromethorphan HBr, USP 20 mg; Guaifenesin, USP 400 mg

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Drug Facts	
Active ingredients (in each 20 ml)	Purposes
Dextromethorphan HBr, USP 20 mg...	Cough suppressant
Guaifenesin, USP 400 mg	Expectorant
Uses	
<ul style="list-style-type: none">temporarily relieves cough due to minor throat and bronchial irritation as may occur with a coldhelps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes	
Warnings	
<p>Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p>	
Ask a doctor before use if you have	
<ul style="list-style-type: none">cough that occurs with too much phlegm (mucus)cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema	
Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.	

Children's Robitussin Elderberry Cough + Chest Congestion DM

Dextromethorphan HBr, USP 20 mg; Guaifenesin, USP 400 mg

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product ■ ml = milliliter

age	dose
children under 4 years	do not use
children 4 to under 6 years	5 ml every 4 hours
children 6 to under 12 years	10 ml every 4 hours
adults and children 12 years and over	20 ml every 4 hours

Other information

- each 20 ml contains: **potassium 5 mg, sodium 20 mg**
- store at 20-25°C (68-77°F). Do not refrigerate.

Children's Robitussin Elderberry Cough + Chest Congestion DM

Dextromethorphan HBr, USP 20 mg; Guaifenesin, USP 400 mg

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Inactive ingredients

anhydrous citric acid, black elderberry juice concentrate (for flavor), carboxymethylcellulose sodium, glycerin, liquid glucose, maltodextrin, natural and artificial flavors, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium gluconate, sucralose, xanthan gum, zinc gluconate

Questions or comments? call weekdays from
9 AM to 5 PM EST at **1-800-245-1040**