

Do Not Use if breakable ring is separated or missing.
Packaged with Tamper-Evident bottle cap.

Robitussin

Drug Facts

Active ingredients **Purposes**
(in each 10 mL)
Dextromethorphan HBr, USP 15 mg... Cough suppressant
Doxylamine succinate, USP 6.25 mg.....Antihistamine

Uses
■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
■ runny nose ■ sneezing ■ itchy, watery eyes
■ itching of the nose or throat
■ controls the impulse to cough to help you sleep

Warnings
Do not use
■ to sedate a child or to make a child sleepy
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ trouble urinating due to an enlarged prostate gland
■ glaucoma
■ a cough that occurs with too much phlegm (mucus)
■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product
■ do not use more than directed
■ marked drowsiness may occur
■ avoid alcoholic drinks
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ be careful when driving a motor vehicle or operating machinery
■ excitability may occur, especially in children

Drug Facts (continued)
Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ measure only with dosing cup provided
■ keep dosing cup with product ■ mL = milliliter
■ do not take more than 4 doses in any 24-hour period

age	dose
children under 6 years	do not use
children 6 to under 12 years	10 mL every 6 hours
adults and children 12 years and older	20 mL every 6 hours

Other information
■ each 10 mL contains: sodium 10 mg
■ store at 20-25°C (68-77°F)

Inactive ingredients
anhydrous citric acid, blueberry juice concentrate, carboxymethylcellulose sodium, glycerin, lactic acid, natural and artificial flavors, natural grade A honey, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium gluconate, sucralose, xanthan gum, zinc gluconate

Questions or comments? call weekdays from 9 AM to 5 PM EST at 1-800-245-1040

Dist. by: Haleon, Warren, NJ 07059
©2024 Haleon or licensor.
Trademarks owned or licensed by Haleon.
For most recent product information, visit www.robitussin.com
Pat. Info. www.productpats.com Made in Canada
62000000211561

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Children's

Robitussin



Nighttime Cough **DM**

Children's

Children's

Robitussin Robitussin



For Ages 6+

Nighttime Cough **DM**

DEXTROMETHORPHAN HBr
(Cough Suppressant)
DOXYLAMINE SUCCINATE (Antihistamine)


ALCOHOL FREE
LONG-ACTING
Relieves:
✓ Cough up to 8 hours
✓ Runny nose

This dynamic duo combines the great taste of natural honey with the effective cough relief of Robitussin® you trust to help your child feel better.

Made with **Real Honey**



Gluten Free
No Artificial Colors



Taste the **Real Honey**
TRUE SOURCE CERTIFIED HONEY
4 FL OZ (118 mL)

85% FPO
3 0031 8762 12 7