

ADULT

**Robitussin**

MAXIMUM STRENGTH

**SEVERE  
MULTI-SYMPOM  
Cough Cold + Flu**



HALEON

Do Not Use if breakable ring on bottle cap is separated or missing.

ADULT

**Robitussin Robitussin**

MAXIMUM STRENGTH

**SEVERE  
MULTI-SYMPOM  
Cough Cold + Flu**

**ACETAMINOPHEN (Pain Reliever/Fever Reducer)  
DEXTROMETHORPHAN HBr (Cough Suppressant)  
GUAIFENESIN (Expectorant)**

- ✓ Cough, Sore Throat
- ✓ Head & Body Aches, Fever
- ✓ Chest Congestion
- ✓ Thins & Loosens Mucus



**FAST, POWERFUL  
multi-symptom relief**

**CF  
MAX**

For Ages 12 & Over

8 FL OZ (237 mL)

**STRONG  
SOOTHING  
ACTION**  
and powerful  
relief for your  
worst symptoms



Also available:



**PARENTS:**  
Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

Distributed by: Haleon, Warren, NJ 07059  
For most recent product information, visit [www.robitussin.com](http://www.robitussin.com)  
Trademarks owned or licensed by Haleon.  
Pat. Info [www.productpats.com](http://www.productpats.com)  
©2023 Haleon or licenser. Made in Canada

**Drug Facts**

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg	Pain reliever/Fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant

**Uses**

- temporarily relieves these symptoms occurring with a cold or flu:
  - cough due to minor throat and bronchial irritation
  - minor aches and pains
  - sore throat
  - headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in any 24-hour period, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have**

- liver disease
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking any other pain reliever/fever reducer

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

- do not take more than 6 doses in any 24-hour period
- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage.
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

**Other information**

- each 20 mL contains: sodium 14 mg
- store at 20-25°C (68-77°F)

**Inactive ingredients** anhydrous citric acid, edetate disodium, FD&C red no. 40, glycerin, menthol, natural and artificial flavors, polyethylene glycol, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, triacetin, xanthan gum

**Questions or comments?** call weekdays from 8 AM to 6 PM EST at 1-800-245-1040

Maximum Strength claim based on maximum levels of active ingredients per 4 hour dose.

62000000207716



3 00312 09467 5