

Packaged with Tamper-Evident ring is separated or missing.
Do Not Use

Children's

Robitussin

AGES 6 & OVER

Drug Facts

Active ingredients (in each 10 mL)

Chlorpheniramine maleate, USP 2 mg Anthistamine
Dextromethorphan HBr, USP 15 mg Cough suppressant

Purposes

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat

Warnings

Do not use

- to sedate a child or to make a child sleepy
 - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.
- If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- a cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- do not use more than directed
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Drug Facts (continued)

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- measure only with dosing cup provided
 - keep dosing cup with product
 - do not take more than 4 doses in any 24-hour period
- mL = milliliter

age	dose
children under 6 years	do not use
children 6 to under 12 years	10 mL every 6 hours
adults and children 12 years and over	20 mL every 6 hours

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)

Inactive ingredients: anhydrous citric acid, FD&C red no. 40, glycerin, lactic acid, natural and artificial flavors, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments? call weekdays from 8 AM to 6 PM EST at 1-800-245-1040

Distributed by: Haleon, Warren, NJ 07059
For most recent product information, visit www.robitussin.com
Trademarks owned or licensed by Haleon.
©2024 Haleon or licensor.
Pat. Info www.productpats.com
Made in Canada



62000000214571



Children's

Robitussin

AGES 6 & OVER

Cough & Cold

Long-Acting

Children's

Robitussin

AGES 6 & OVER

Cough & Cold

Long-Acting

CHLORPHENIRAMINE MALEATE (Antihistamine)
DEXTROMETHORPHAN HBr (Cough Suppressant)

Relieves:

- ✓ Cough Up to 8 Hours
- ✓ Runny Nose

Alcohol-Free



fruit punch
flavor

4 FL OZ
(118 mL)

Children's Robitussin liquid is specially formulated to provide soothing action, control your child's cough plus relieve other cold symptoms.



PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org