

Robitussin Maximum Strength Cough & Chest Congestion DM

Dextromethorphan HBr, USP 10 mg, Guaifenesin, USP 200 mg

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Drug Facts

Active ingredients (in each liquid-filled capsule) Purposes

Dextromethorphan HBr, USP 10 mg Cough suppressant
Guaifenesin, USP 200 mg Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Robitussin Maximum Strength Cough & Chest Congestion DM

Dextromethorphan HBr, USP 10 mg, Guaifenesin, USP 200 mg

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Directions

- do not take more than 12 capsules in any 24-hour period
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	2 capsules every 4 hours
children under 12 years	do not use

Other information

- store at 20-25°C (68-77°F).
- avoid excessive heat above 40°C (104°F).

Inactive ingredients

FD&C Red No. 40, gelatin, glycerin, light mineral oil, pharmaceutical ink, polyethylene glycol, providone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments?

Call weekdays from 9 AM to 5 PM EST at **1-800-762-4675**