

# Robitussin Maximum Strength Nighttime Cough DM

## Dextromethorphan HBr, Doxylamine Succinate

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

<b>Drug Facts</b>	
<b>Active ingredients (in each 20 ml)</b>	<b>Purposes</b>
Dextromethorphan HBr 30 mg .....	Cough Suppressant
Doxylamine Succinate, USP 12.5 mg.....	Antihistamine
<b>Uses</b>	
<ul style="list-style-type: none"><li>■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold</li><li>■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:<ul style="list-style-type: none"><li>■ runny nose</li><li>■ sneezing</li><li>■ itchy, watery eyes</li><li>■ itching of the nose or throat</li></ul></li><li>■ controls the impulse to cough to help you sleep</li></ul>	
<b>Warnings</b>	
<b>Do not use</b>	
<ul style="list-style-type: none"><li>■ to sedate a child or to make a child sleepy</li><li>■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li></ul>	
<b>Ask a doctor before use if you have</b>	
<ul style="list-style-type: none"><li>■ trouble urinating due to an enlarged prostate gland</li><li>■ glaucoma</li><li>■ a cough that occurs with too much phlegm (mucus)</li><li>■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema</li></ul>	
<b>Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers</b>	

# Robitussin Maximum Strength Nighttime Cough DM

## Dextromethorphan HBr, Doxylamine Succinate

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

### When using this product

- do not use more than directed
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Stop use and ask a doctor if** cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- measure only with dosing cup provided
- keep dosing cup with product
- ml = milliliter
- do not take more than 4 doses in any 24-hour period
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 ml every 6 hours
children under 12 years	do not use

### Other information

- each 20 ml contains: sodium 14 mg
- store at 20-25°C (68-77°F)

### Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, FD&C blue no. 1, FD&C red no. 40, glycerin, liquid glucose, menthol, natural and artificial flavors, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose, triacetin, xanthan gum

### Questions or comments?

call weekdays from 9 AM to 5 PM EST at **1-800-762-4675**