

Robitussin Honey Severe Cough, Flu + Sore Throat CF Max Nighttime

Acetaminophen (Pain reliever/Fever reducer), Diphenhydramine HCl
(Antihistamine/Cough suppressant)

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Drug Facts	
Active ingredients (in each 20 ml)	Purposes
Acetaminophen, USP 650 mg	Pain reliever/Fever reducer
Diphenhydramine HCl, USP 25 mg	Antihistamine/Cough suppressant
Uses	
<ul style="list-style-type: none"> ■ temporarily relieves these symptoms occurring with a cold or flu, hay fever, or other upper respiratory allergies: <ul style="list-style-type: none"> ■ cough due to minor throat and bronchial irritation ■ minor aches and pains ■ sore throat pain ■ headache ■ runny nose, sneezing, itchy watery eyes ■ itching of the nose and throat ■ temporarily relieves your cough to help you sleep ■ temporarily reduces fever 	
Warnings	
<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> ■ more than 6 doses in any 24-hour period, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product 	
<p>Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> ■ skin reddening ■ blisters ■ rash <p>If a skin reaction occurs, stop use and seek medical help right away.</p>	
<p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>	
Do not use	
<ul style="list-style-type: none"> ■ to sedate a child or to make a child sleepy ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ with any other product containing diphenhydramine, even one used on skin 	

Robitussin Honey Severe Cough, Flu + Sore Throat CF Max Nighttime

Acetaminophen (Pain reliever/Fever reducer), Diphenhydramine HCl
(Antihistamine/Cough suppressant)

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Ask a doctor before use if you have

- liver disease
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, emphysema or chronic bronchitis
- if you have glaucoma or difficulty in urination due to enlargement of prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin
- taking any other pain reliever/fever reducer

When using this product

- **do not use more than directed**
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

Robitussin Honey Severe Cough, Flu + Sore Throat CF Max Nighttime

Acetaminophen (Pain reliever/Fever reducer), Diphenhydramine HCl
(Antihistamine/Cough suppressant)

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Directions

- do not take more than 6 doses in any 24-hour period
- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage.
- measure only with dosing cup provided
- keep dosing cup with product
- ml = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 ml every 4 hours
children under 12 years	do not use

Other information

- each 20 ml contains: **sodium 22 mg**
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, glycerin, natural & artificial flavors, natural grade A honey, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium gluconate, sucralose, xanthan gum

Questions or comments?

call weekdays from 9 AM to 5 PM EST at **1-800-762-4675**