

Children's Robitussin 12 Hour Cough Relief

Dextromethorphan Polistirex Extended-Release Oral Suspension

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Drug Facts

Active ingredient (in each 5 mL)**Purpose**

Dextromethorphan polistirex equivalent to 30 mg
dextromethorphan hydrobromide, USP Cough suppressant

Uses temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

Children's Robitussin 12 Hour Cough Relief

Dextromethorphan Polistirex Extended-Release Oral Suspension

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Directions

- **shake bottle well before use**
- measure only with dosing cup provided. Do not use dosing cup with other products
- dose as follows or as directed by doctor
- mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: **sodium 5 mg**
- store at 20-25°C (68-77°F)
- dosing cup provided

Inactive ingredients (Orange flavor)

D&C red no. 30, D&C yellow no. 10, flavor, glycerin, high fructose corn syrup, methylparaben, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Inactive ingredients (Grape flavor)

D&C red no. 30, FD&C blue no. 1, flavor, glycerin, high fructose corn syrup, methylparaben, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions? call weekdays from 9 AM to 5 PM EST at **1-800-762-4675**. You may also report side effects to this number.