

# Robitussin

## Lingering Cold Long-Acting CoughGels

Dextromethorphan HBr, USP

This is the most current labeling information, and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

### **Drug Facts**

<b>Active ingredient (in each liquid-filled capsule)</b>	<b>Purpose</b>
Dextromethorphan HBr, USP 15 mg .....	Cough suppressant

**Use** temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold.

### **Warnings**

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- a cough that occurs with too much phlegm (mucus)
- a cough that lasts or is chronic as occurs with smoking, asthma, or emphysema

**Stop use and ask a doctor if** cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

# Robitussin

## Lingering Cold Long-Acting CoughGels

Dextromethorphan HBr, USP

This is the most current labeling information, and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

### **Directions**

- do not take more than 8 capsules in any 24-hour period
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	take 2 capsules every 6 to 8 hours, as needed
children under 12 years	do not use

### **Other information**

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)
- protect from light

### **Inactive ingredients**

FD&C blue no. 1, FD&C red no. 40, fractionated coconut oil, gelatin, glycerin, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propyl gallate, propylene glycol, purified water, sorbitol, sorbitol anhydrides

### **Questions or comments?**

Call weekdays from 9 AM to 5 PM EST at **1-800-762-4675**.

## Document Approval Record

**Document Name:**

LBL-00000183 WEB READY - ROBITUSSIN LINGERING COLD LON  
G-ACTING COUGHGELS (Version 13)

**Document Title:**

LBL-00000183 WEB READY - ROBITUSSIN LINGERING COLD LON  
G-ACTING COUGHGELS (Version 13)

**Signed By:****Date(GMT)****Signing Capacity**

Strotz, Carl

21-Sep-2017 19:47:48

Final Approval

Mazurets, Natalie

21-Sep-2017 19:55:14

Final Approval

Hillebrecht, Christy

21-Sep-2017 19:56:49

Final Approval