

Children's Robitussin Cough & Cold CF

Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Drug Facts

Active ingredients (in each 10 ml)

Dextromethorphan HBr, USP 10 mg	Cough suppressant
Guaifenesin, USP 100 mg	Expectorant
Phenylephrine HCl, USP 5 mg	Nasal decongestant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

When using this product do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Children's Robitussin Cough & Cold CF

Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Directions

- measure only with dosing cup provided
- keep dosing cup with product
- ml = milliliter
- do not take more than 6 doses in any 24-hour period

age	dose
children under 4 years	do not use
children 4 to under 6 years	5 ml every 4 hours
children 6 to under 12 years	10 ml every 4 hours
adults and children 12 years and over	20 ml every 4 hours

Other information

- each 10 ml contains: sodium 7 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, artificial flavor, FD&C red no. 40, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments? call weekdays from 9 AM to 5 PM EST at
1-800-762-4675