

Children's Robitussin Cough & Chest Congestion DM

Dextromethorphan HBr 5 mg, Guaifenesin 100 mg

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Drug Facts	
Active ingredients (in each 5 ml)	Purposes
Dextromethorphan HBr, USP 5 mg	Cough suppressant
Guaifenesin, USP 100 mg	Expectorant
Uses	
<ul style="list-style-type: none">■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes	
Warnings	
<p>Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p>	
Ask a doctor before use if you have	
<ul style="list-style-type: none">■ cough that occurs with too much phlegm (mucus)■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema	
<p>Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.</p>	
<p>If pregnant or breast-feeding, ask a health professional before use.</p>	
<p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	

Children's Robitussin Cough & Chest Congestion DM

Dextromethorphan HBr 5 mg, Guaifenesin 100 mg

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Directions

- measure only with dosing cup provided
- keep dosing cup with product
- ml = milliliter
- do not take more than 6 doses in any 24-hour period

age	dose
children under 4 years	do not use
children 4 to under 6 years	5 ml every 4 hours
children 6 to under 12 years	10 ml every 4 hours
adults and children 12 years and over	20 ml every 4 hours

Other information

- each 5 ml contains: sodium 3 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid, artificial flavor, FD&C blue no. 1, FD&C red no. 40, glycerin, natural flavor, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments? call weekdays from 9 AM to 5 PM EST at 1-800-762-4675