

Robitussin Maximum Strength 12 Hour Cough & Mucus Relief Extended-Release Tablets

Guaifenesin 1200 mg, Dextromethorphan Hydrobromide 60 mg

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Drug Facts

Active ingredients (in each extended-release tablet)	Purposes
Dextromethorphan HBr 60 mg.....	Cough suppressant
Guaifenesin 1200 mg.....	Expectorant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep

Warnings

Do not use

- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

When using this product

- do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Robitussin Maximum Strength 12 Hour Cough & Mucus Relief Extended-Release Tablets

Guaifenesin 1200 mg, Dextromethorphan Hydrobromide 60 mg

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

Other information

- **each tablet contains:** magnesium 25 mg
- store between 20-25°C (68-77°F)

Inactive ingredients

carbomer homopolymer type B, copovidone, D&C yellow #10 aluminum lake, hypromellose, magnesium hydroxide, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions? Call weekdays from 9 AM to 5 PM EST at **1-800-762-4675**